

**DANISH METEOROLOGICAL
INSTITUTE**

— TECHNICAL REPORT —

02-24

**Verification of DMI wave forecasts
2nd quarter of 2002**

Jacob Woge Nielsen



Copenhagen 2002

ISSN 0906-897X (trykt)
ISSN 1399-1388 (online)

Verification of DMI wave forecasts

2nd quarter of 2002

Jacob Woge Nielsen
e-mail: jw@dmi.dk

The Danish Meteorological Institute, Copenhagen, Denmark

Contents

| | | |
|------------|--|-----------|
| 1 | Introduction with Key Numbers | 3 |
| 2 | DMI-WAM | 4 |
| 2.1 | Physical model | 4 |
| 2.2 | Model set-up | 4 |
| 2.3 | Weather model | 5 |
| 3 | Wave data | 6 |
| 3.1 | Forecasts | 6 |
| 3.2 | Observations | 6 |
| 4 | Error measures | 9 |
| 5 | Results | 10 |
| 5.1 | Significant wave height | 10 |
| 5.2 | Extreme wave height | 13 |
| 5.3 | Mean wave period | 14 |
| 5.4 | Dominant wave period | 15 |
| 5.5 | Mean wave direction | 16 |
| 6 | Conclusion | 17 |
| 7 | Appendix | 18 |
| 7.1 | Wave recorders | 18 |
| 7.2 | Observed wave statistics | 19 |
| 7.3 | Wave height vs. wave period | 21 |
| 7.4 | Wave forecast statistics | 22 |
| 7.5 | Significant wave height station plots | 25 |
| 7.6 | Mean wave period station plots | 56 |
| 7.7 | Dominant wave period station plots | 61 |

1 Introduction with Key Numbers

We analyse the quality of wave forecasts valid for the 2nd quarter of 2002, produced by *DMI-WAM* - DMI's operational set-up of the 3rd generation wave model *WAM Cycle4*.

The significant wave height H_s is of primary interest, but other wave parameters (wave period, direction, swell parameters) are examined where the data material is adequate.

Standard error measures (bias, rms error, ..) are calculated both as a function of forecast range and of wave height. Special statistics are done for extreme waves. Grand averages are calculated as mean values over all stations, over all ranges, and for separate geographical regions.

All model results are forecasts. This means that errors in the parameters do not necessarily imply errors of the model, but may reflect errors in the meteorological forcing data and initial conditions.

DMI has produced short-range operational wave forecasts since 1999. A pre-operational validation study was carried out in 1999 [7], a combined wave-wind validation in 2000 for a 14 month hindcast period [6], and a verification pilot study in 2002 [5]. Previous report(s) are [4].

Outline: Ch. 2 briefly describes the DMI wave model set-up, ch. 3 lines out the data material, and in ch. 4 we define the statistical error measures used to describe the forecast quality. Ch. 5 presents and discusses the results. Ch. 6 concludes the work. Comprehensive results for each station are found in the Appendix. References and lists of figures/tables are found at the end of the report.

For convenience, the Table below shows *Key Numbers* pertaining to the full model system. Please refer to the Results section (ch. 5) for a detailed explanation and discussion.

| Parameter | H_s | T_{02} | T_p | θ_w |
|----------------|-------|----------|-------|------------|
| bias | -2cm | 1.1s | 1.6s | 7° |
| relative bias | 3% | 31% | 26% | |
| rms error | 42cm | 1.4s | 2.8s | |
| st.dev | | | | 47° |
| scatter index | 0.34 | 0.40 | 0.46 | |
| corr.coeff. | 0.86 | 0.48 | 0.52 | 0.85 |
| peak bias | -21cm | | | |
| rel. peak bias | -4% | | | |

Table 1. Key numbers

2 DMI-WAM

DMI runs an operational wave forecasting service for Danish waters since 1999, using the 3rd generation wave model *WAM Cycle4* (described in detail in [1], [3]) forced by DMI's numerical weather prediction model *HIRLAM*. In 2002, the geographical model domain includes a large part of the North Atlantic, the North Sea and Baltic Sea, and the Mediterranean. The DMI-WAM model set-up is described in detail below.

2.1 Physical model

WAM Cycle4 solves the spectral wave equation

$$\frac{\partial F}{\partial t} + \vec{c} \cdot \vec{\nabla} F = S_{in} + S_{nl} + S_{ds} + S_{cu} + S_{bf}$$

where $F(f, \theta; \vec{x}, t)$ is spectral wave energy density, depending on wave frequency, wave direction, position and time; $c(f, \theta)$ is the wave group speed; S_{in} is wind energy input; S_{nl} is non-linear wave-wave interaction; S_{ds} is wave energy dissipation through wave breaking (white capping); S_{cu} is wave-current interaction; S_{bf} is interaction with the sea bed through friction and wave refraction. DMI-WAM still lacks current data ($S_{cu}=0$) and information about sea ice.

2.2 Model set-up

DMI provides wave forecasts in three geographical domains as shown below:




Figure 1. DMI wave models. North Atlantic, North Sea-Baltic, and Mediterranean model

The model open boundaries are chosen as follows. The coarse grid North Atlantic model uses the JONSWAP wind-sea spectrum (see [1], [2]). The fine grid North Sea - Baltic model is nested into the North Atlantic model, and uses spatially interpolated wave spectra calculated by that model. The Mediterranean is treated as a closed basin, assuming no wave energy exchange with the Atlantic or the Black Sea. Please refer to Table 2 for a model set-up summary.

The wave forecasting system was coldstarted using developed sea. Subsequent model runs are initialised using the sea state at analysis time, calculated by the previous run.

| Model | North Atlantic | North Sea - Baltic | Mediterranean |
|------------------|----------------|--------------------|---------------|
| Space res. | 30' | 10' | 10' |
| Time step | 4 min | 4 min | 2 min |
| Frequencies | 25 | 25 | 25 |
| Direction resol. | 30° | 30° | 30° |
| Forcing model(s) | Hirlam G | Hirlam E | Hirlam E+G |
| - resolution of | 0.45° | 0.15° | 0.15°/0.45° |
| Longitudes | 69°W-30°E | 20°W-30°E | 6°W-46°E |
| Latitudes | 30°N-75°N | 36°N-68°N | 30.5°N-46°N |
| Open boundaries | JONSWAP | Nested | Closed basin |
| Forecast range | 54 h | 54 h | 54 h |
| Output time step | 1 h | 1 h | 1 h |
| Schedule | 4x daily | 4x daily | 4x daily |

Table 2. DMI-WAM set-up. The wave model frequencies range from 0.04177 Hz to 0.41145 Hz in 10% steps. The Mediterranean model patches Hirlam E+G to get maximum resolution. The directional resolution is 30° (in previous reports erroneously stated as 15°). Changes to the model set-up usually require a new coldstart.

2.3 Weather model

The forcing models are the DMI limited area numerical weather prediction models Hirlam-E and Hirlam-G. Both are currently being used in the DMI weather forecasting service. The G model covers a larger area than the E model, but in coarser spatial resolution (45km vs. 15km on a rotated latitude-longitude grid). The wind vector at 10m height is interpolated linearly in time and space to match the spherical wave model grids.




Figure 2. DMI Hirlam. The outermost box is the G model, the box covering most of Europe is the E model.

3 Wave data

The verification data consists of operational DMI-WAM wave forecasts, and wave observations from a number of fixed positions (buoys and platforms).

3.1 Forecasts

Wave parameters output from DMI-WAM are shown in Table 3.

| DMI-WAM output | |
|----------------|-------------------------|
| H_s | Significant wave height |
| H_{sw} | Height of swell |
| T_{02} | Mean wave period |
| T_p | Dominant wave period |
| T_{sw} | Swell mean period |
| θ_w | Mean wave direction |
| θ_{sw} | Swell direction |

Table 3. DMI-WAM wave parameters

The forecasts are stored as hourly maps in model resolution. Time series for each station are sampled using nearest model grid point. This is done for each analysis and each parameter. During the 2nd quarter of 2002, 364 forecasts were produced, of which 355 were archived successfully on tape.

Most wave parameters are obtained by a suitable integral of the wave energy spectrum F . In contrast, the dominant wave period T_p is the discretized model frequency (inverse) containing the highest energy, picked from the 25 predefined values. In cases with two competing wave energy maxima (wind sea and swell), T_p may flip between the two, as first one, then the other has the highest energy. This means that T_p is a non-smooth function of time or space, sampled with low accuracy.

3.2 Observations

The wave recorder positions are shown in Fig. 3 below. A total of 30 stations that record more or less regularly are selected for verification. Comprehensive station information is found in Appendix 7.1, Table 11.

Wave data is obtained from a number of sources, as indicated in table 4. SMHI, KDI and NCMR data are kindly provided by each agency in question. NDBC data is retrieved via the GTS. Table 5 shows the number of stations for each wave parameter, and for each of 8 geographical domains.

| Wave Data providers | |
|---------------------|--|
| NDBC | National Data Buoy Center (UK) |
| SMHI | Swedish Meteorological Institute |
| KDI | Danish Coastal Authority |
| NCMR | National Center for Marine Research (Greece) |

Table 4. Wave data providers.




Figure 3. Wave recorder locations.

| Parameter | H_s | T_{02} | T_p | θ_w | H_m |
|-----------------------|-------|----------|-------|------------|-------|
| Atlantic | 8 | - | 8 | - | - |
| Scotland-Faroe | 1 | - | 1 | - | - |
| Irish Sea | 2 | - | 2 | - | - |
| Br. Channel | 4 | - | 4 | - | - |
| North Sea | 5 | - | 4 | - | - |
| Danish West Coast | 4 | 4 | 4 | 4 | 4 |
| Kattegat - Baltic Sea | 3 | 3 | - | - | 3 |
| Mediterranean | 3 | 1 | 3 | 1 | - |
| Total | 30 | 8 | 26 | 5 | 7 |

Table 5. Number of wave stations in each domain, and for each wave parameter. Maximum wave height H_m , recorded at Baltic and Danish West Coast stations only, is forecasted by the wave model. Swell parameters are not recorded at any fixed location.

The mean data coverage is 83% (see Fig. 4 for missing data). The sampling rate is 1 hour, with 9 stations (cf. Table 11) sampling every 3 hours. At these stations, we consider steady 3 hour sampling as full data coverage.

The sampling accuracy is H_s :0.1m, T_p :1s, T_{02} :0.01s, θ_w : 1° . Two buoys (62101, 62301) use low 0.5m H_s accuracy. Four buoys (Danish West Coast) use variable T_p accuracy of 0.1-0.7s, while one (Athos) uses high 0.01s T_p accuracy. Statistics on the observed wave parameters, and wave period vs. wave height plots, are shown in Appendices 7.2 and 7.3.




Figure 4. Missing data. St. 20044, 24023, 25077 and 25138 sample every 3rd hour, but revert to 1h sampling in storm situations. St. 61001, 61002, 62117, 62414 and Athos sample every 3rd hour.

4 Error measures

Model errors are calculated using a 3-d residual matrix, built from all available observations and forecasts. With the general formula $\text{residual} = \text{forecast} - \text{observation}$ the matrix reads (brackets indicate a dependency)

$$\text{residual}(\text{station}, \text{analysis}, \text{range})$$

with the number of stations depending on the parameter in question (cf. Table 5), analysis every 6 hours, and forecasts ranging from 1-54 hours in 1 hour steps. With 30 stations, the matrix has 589680 elements. By averaging the residual over all analyses, we get the model bias or mean error:

$$\text{bias}(\text{station}, \text{range})$$

Further linear averaging gives the bias for each forecast range (averaged over all stations), for each station (averaged over the full forecast range), and as a grand average.

$$\begin{aligned} &\text{bias1}(\text{range}) \\ &\text{bias2}(\text{station}) \\ &BIAS \end{aligned}$$

In the same way, the root mean square error $\text{rms}(\text{station}, \text{range})$ is calculated and averaged using the residual squared.

For the wave height only, the bias and rms error are also calculated as a function of wave height. The residual is sorted into observation bins 0.5m wide and averaged for each bin.

$$\begin{aligned} &\text{bias}(\text{station}, \text{obsbin}) \\ &\text{rms}(\text{station}, \text{obsbin}) \end{aligned}$$

Averaging over all stations gives the model error dependency on wave height, calculated both as an absolute value and as a relative error in %.

The scatter index $si = \frac{\text{rms}}{\langle \text{obs} \rangle}$ is obtained by normalising rms with the observed mean value. si may be used to intercompare rms errors at stations with large differences in wave climate. Averaging is done as above.

Correlation coefficients $cc(\text{station}, \text{block})$ are calculated using forecast pseudo time series, established by concatenating forecasts in 6 hour range blocks. This gives coefficients valid for each of the 9 range blocks 1-6, 7-12, .. , 49-54 hours. Range block and station-dependent values ($cc1, cc2$), and a grand average (CC), are calculated.

A special peak bias $pbias(\text{station}, \text{block})$ is calculated using the 5 most extreme events at each station, allowing for a forecast phase error of a few hours. Peak biases are calculated both as absolute and relative values.

5 Results

This section describes wave verification results for the 2nd quarter of 2002, for significant wave height (H_s), mean and dominant wave period (T_{02}, T_p), and mean wave direction (θ_w). In the sections below, we discuss grand averages and regional averages for each wave parameter in turn. Detailed results for each station are found in the Appendix at the end of the report.

5.1 Significant wave height

Table 6 shows bias and relative bias, rms error, scatter index and correlation coefficient, averaged over the full forecast range. Fig. 5 shows the short range (1-6h) scatter diagram. The error estimates are sorted out on 8 geographical regions.

| Parameter Region | #st | bias cm | bias % | rms cm | si | cc |
|---------------------|-----|------------|-----------|-----------|------|------|
| Atlantic | 8 | -18 | -6 | 56 | 0.20 | 0.90 |
| Scotland–Faroe | 1 | -18 | -7 | 49 | 0.19 | 0.90 |
| Irish Sea | 2 | -12 | -9 | 39 | 0.27 | 0.90 |
| Br.Channel | 4 | 30 | 30 | 52 | 0.56 | 0.84 |
| North Sea | 5 | -2 | -1 | 36 | 0.27 | 0.87 |
| Danish West Coast | 4 | 3 | 3 | 27 | 0.33 | 0.86 |
| Kattegat–Baltic | 3 | 8 | 16 | 25 | 0.48 | 0.83 |
| Mediterranean | 3 | -11 | -9 | 41 | 0.47 | 0.73 |
| All Waters | 30 | -2 | 3 | 42 | 0.34 | 0.86 |

Table 6. Significant wave height results




Figure 5. Significant wave height: short range (1-6h) scatter diagram

The wave model has almost no bias on average and an rms error of 0.42 m. Scatter index is low (0.34) and correlation coefficient high (0.86). There is some regional spread. Waves are underpredicted on average (negative bias) at the Atlantic, Irish Sea, Shetland and Mediterranean stations, and overpredicted (positive bias) in the British Channel and in the Baltic Sea. RMS errors range roughly from a quarter to half a metre. The scatter index is well below an acceptance level of 0.4 in most

regions, but exceeds in the British Channel, in the Baltic and in the Mediterranean. Fig. 6 shows the scatter index at each station.




Figure 6. Significant wave height: scatter index

The error dependency on forecast range and on wave height is shown in Fig. 7.

The model bias is almost independent of forecast range, while the rms error/scatter index increase slowly. The rms error is significant already at analysis time since the model is initialised without any use of the observed sea state. Conversely, the correlation coefficient decreases, from 0.89 at short range to 0.80 at long range.

There is a strong dependency on wave height. Very small waves are slightly overestimated, while higher waves are underestimated by up to about 0.5m on average. The rms error increases steadily with increasing wave height. Except for very small waves (less than 1m), the relative bias and rms error only depend weakly on wave height. The relative rms error decreases slowly with wave height, with a standard level of about 20%. For waves >8m the data material is too sparse to produce reliable statistics.

Results on significant wave height for each single station is shown in Table 15, Appendix 7.4. Data sheets are presented in Appendix 7.5.




Figure 7. Significant wave height

5.2 Extreme wave height

The error on extreme waves is calculated by singling out the 5 highest events at each station, and then calculate the forecast error, allowing for a few hours' phase displacement. Table 7 shows peak biases for each of the 8 geographical domains, averaged over all forecast ranges.

| Parameter Region | #st | peak bias | |
|---------------------|-----|-----------|-----|
| | | cm | % |
| Atlantic | 8 | -60 | -10 |
| Scotland–Faroe | 1 | -52 | -6 |
| Irish Sea | 2 | 6 | 0 |
| Br.Channel | 4 | 41 | 12 |
| North Sea | 5 | -8 | 0 |
| Danish West Coast | 4 | -7 | -3 |
| Kattegat–Baltic | 3 | -22 | -11 |
| Mediterranean | 3 | -41 | -13 |
| All Waters | 30 | -21 | -4 |

Table 7. Extreme wave height results

There is a negative peak bias in most regions, with the British Channel as an exception. On average, the system has a small negative peak bias of -21cm or -4%.

The dependency of the peak bias on forecast range is shown in Fig. 8. The peak bias is small at short range (day 1), but increases on day 2 and beyond.




Figure 8. Extreme wave height

Table 16 in Appendix 7.4 shows peak biases for each single station.

5.3 Mean wave period

The mean wave period T_{02} is recorded at 8 stations. Grand averages are shown in Table 8, and a short-range scatter diagram in Fig. 9.

| Parameter Region | #st | bias | | rms | si | cc |
|-------------------|-----|------|----|------|------|------|
| | | sec. | % | sec. | | |
| Danish West Coast | 4 | 1.7 | 47 | 2.2 | 0.60 | 0.40 |
| Kattegat–Baltic | 3 | 0.5 | 13 | 0.7 | 0.20 | 0.59 |
| Mediterranean | 1 | 0.5 | 16 | 0.7 | 0.21 | 0.43 |
| All Waters | 8 | 1.1 | 31 | 1.4 | 0.40 | 0.48 |

Table 8. Mean wave period results

At the Danish West Coast stations T_{02} is overestimated by almost 50%. The reason for this is unresolved. At the stations in the Baltic and the Mediterranean T_{02} is only slightly overestimated. The scatter index at these stations is well below the acceptancy level of 0.4.




Figure 9. Mean wave period, 1-6h range. Left panel: all stations, right panel: Baltic stations only

Results on mean wave period for each station are shown in Table 17, Appendix 7.4. Data sheets are presented in Appendix 7.6.

5.4 Dominant wave period

The dominant (or *peak*) wave period T_p is recorded at 26 stations. Grand averages are shown in Table 9, short-range scatter diagrams in Fig. 10.

| Parameter Region | #st | bias sec. | % | rms sec. | si | cc |
|---------------------|-----|--------------|----|-------------|------|------|
| Atlantic | 8 | 2.7 | 35 | 3.1 | 0.41 | 0.67 |
| Scotland–Faroe | 1 | 2.9 | 42 | 3.4 | 0.50 | 0.59 |
| Irish Sea | 2 | 2.7 | 51 | 3.6 | 0.67 | 0.56 |
| Br.Channel | 4 | 0.4 | 13 | 2.9 | 0.50 | 0.37 |
| North Sea | 4 | 1.8 | 38 | 2.6 | 0.54 | 0.35 |
| Danish West Coast | 4 | 0.6 | 10 | 2.6 | 0.45 | 0.57 |
| Mediterranean | 3 | -0.1 | -1 | 1.4 | 0.28 | 0.40 |
| All Waters | 26 | 1.6 | 26 | 2.8 | 0.46 | 0.52 |

Table 9. Dominant wave period results

T_p shows very bad verification results due partly to low recording and forecasting accuracy, and partly to the non-smoothness of the series, with T_p shifting abruptly between a high and a low period peak. Even when the wave spectrum is rather well predicted, a small error in the shape of the spectrum may lead to very large T_p errors in situations with a two-peaked spectrum (swell and wind sea).




Figure 10. Dominant wave period, 1-6h range. Left panel: all stations, right panel: Mediterranean stations only

Results on dominant wave period for each stations is found in Table 18, Appendix 7.4. Data sheets in Appendix 7.7.

5.5 Mean wave direction

The mean wave direction θ_w is recorded at 5 stations. The results are presented in Table 10 and the scatter diagram in Fig. 11.

| Parameter Region | #st | bias deg. | std deg. | cc |
|-------------------|-----|-----------|----------|------|
| Danish West Coast | 4 | 5 | 47 | 0.84 |
| Mediterranean | 1 | 14 | 49 | 0.89 |
| All Waters | 5 | 7 | 47 | 0.85 |

Table 10. Mean wave direction results

The mean wave direction predictions fit the observations rather well, with almost no bias and a high correlation coefficient.




Figure 11. Mean wave direction, 1-6h range.

Results on the mean wave direction for each stations is found in Table 19, Appendix 7.4. Data sheets are not shown.

6 Conclusion

DMI wave forecasts valid for the 2nd quarter of 2002 are verified, using wave data from 30 buoys. A Key Number Table is put right after the Introduction in the beginning of the report (Table 1).

Main conclusions are:

- The significant wave height H_s and mean wave direction θ_w are well predicted, but we have some problems predicting wave period
- There is a large regional spread in forecast quality
- The H_s model error depends mainly on wave height. Extreme waves are usually underestimated
- The forecast quality decreases slowly with increasing forecast range

The *significant wave height* is recorded at all stations. The error distribution is examined in terms of forecast range, as a function of observed wave height, and for separate geographical regions.

The observation-forecast correlation is high, 0.86 on average. This falls off a bit as the prediction range increases.

The bias is very small on average, but there is a large geographical spread, and some dependency on wave height. Low waves are overpredicted, while the highest waves are underestimated by about 5%. The bias does not depend on forecast range.

The average rms error is 0.42m, increasing gradually with forecast range. Except for situations with almost calm sea, waves have a relative rms error of about 20%. An average scatter index SI=0.34 is acceptable. 9 stations have si>0.4 (sometimes used as an acceptance level) due to low recording accuracy or small average wave height.

Extreme waves are underestimated in most regions, with an average -4% peak bias. An exception is the British Channel, where the highest waves are overestimated by about 12%. The negative bias increases on day 2 and beyond.

Two types of wave period are recorded; mean and dominant (peak) wave period.

The *mean wave period* is recorded at 8 locations, half of which have a data interpretation problem. At the remaining 4 stations the model overestimates the mean wave period by roughly half a second, with a scatter index of 0.20 and a correlation coefficient of 0.51.

Dominant wave period predictions are not good. This is a data problem; a well predicted wave spectrum does not guarantee a correct dominant wave period in situations with two spectral maxima. Also, most stations sample only with 1s accuracy and so does the model; this in itself leads to large error measures. Only the Mediterranean stations show good results, with no bias and a scatter index of 0.28.

Mean wave direction predictions have a small bias, a standard deviation of 47°, and a high correlation coefficient.

Swell parameters are not recorded at any of the fixed positions. A few record *maximum wave height* but this is not predicted by the wave model.

7 Appendix

This Appendix contains a wave recorder station table (below), observation statistics tables, forecast statistics tables, wave height/period plots, and a plot sheet for each station and each parameter ($H_s, T_{02}, T_p, \theta_w$), arranged sequentially according to the station table.

7.1 Wave recorders

| Station ID | Agency | Region | lat. | lon. | δt | parameters |
|------------|--------|---------------|--------|--------|------------|-----------------------------------|
| almag | SMHI | Baltic | 59.15N | 19.13E | 1h | H_s, H_m, T_{02} |
| oland | SMHI | Baltic | 56.07N | 16.68E | 1h | H_s, H_m, T_{02} |
| truba | SMHI | Baltic | 57.60N | 11.63E | 1h | H_s, H_m, T_{02} |
| 20044 | KDI | D. West Coast | 57.58N | 9.41E | 3h | $H_s, H_m, T_{02}, T_p, \theta_w$ |
| 24023 | KDI | D. West Coast | 56.47N | 8.06E | 3h | $H_s, H_m, T_{02}, T_p, \theta_w$ |
| 25077 | KDI | D. West Coast | 55.81N | 7.94E | 3h | $H_s, H_m, T_{02}, T_p, \theta_w$ |
| 25138 | KDI | D. West Coast | 55.35N | 8.23E | 3h | $H_s, H_m, T_{02}, T_p, \theta_w$ |
| 61001 | NDBC | Mediterranean | 43.4N | 7.8W | 3h | H_s, T_p |
| 61002 | NDBC | Mediterranean | 42.1N | 4.7W | 3h | H_s, T_p |
| 62001 | NDBC | Atlantic | 45.2N | 5.0W | 1h | H_s, T_p |
| 62026 | NDBC | North Sea | 55.3N | 1.1E | 1h | H_s, T_p |
| 62029 | NDBC | Atlantic | 48.7N | 12.4W | 1h | H_s, T_p |
| 62101 | NDBC | B.Channel | 50.6N | 2.7W | 1h | H_s, T_p |
| 62103 | NDBC | B.Channel | 49.9N | 2.9W | 1h | H_s, T_p |
| 62105 | NDBC | Atlantic | 54.9N | 12.6W | 1h | H_s, T_p |
| 62106 | NDBC | Atlantic | 57.0N | 9.9W | 1h | H_s, T_p |
| 62107 | NDBC | Atlantic | 50.1N | 6.1W | 1h | H_s, T_p |
| 62108 | NDBC | Atlantic | 53.5N | 19.5W | 1h | H_s, T_p |
| 62109 | NDBC | North Sea | 57.0N | 0.0 | 1h | H_s, T_p |
| 62117 | NDBC | North Sea | 57.9N | 0.0 | 3h | H_s, T_p |
| 62145 | NDBC | North Sea | 53.1N | 2.8E | 1h | H_s, T_p |
| 62163 | NDBC | Atlantic | 47.5N | 8.5W | 1h | H_s, T_p |
| 62301 | NDBC | Irish Sea | 52.3N | 4.5W | 1h | H_s, T_p |
| 62303 | NDBC | Irish Sea | 51.6N | 5.1W | 1h | H_s, T_p |
| 62304 | NDBC | B.Channel | 51.1N | 1.8E | 1h | H_s, T_p |
| 62305 | NDBC | B.Channel | 50.4N | 0.0 | 1h | H_s, T_p |
| 62414 | NDBC | North Sea | 53.8N | 2.9E | 3h | H_s |
| 64045 | NDBC | Atlantic | 59.1N | 11.4W | 1h | H_s, T_p |
| 64046 | NDBC | Scotland | 60.7N | 4.5W | 1h | H_s, T_p |
| athos | NCMR | Mediterranean | 39.96N | 24.72E | 3h | $H_s, T_{02}, T_p, \theta_w$ |

Table 11. Wave stations. Station name/number, driving agency, position, and wave parameters. SMHI=Swedish Meteorological Institute, NDBC=National Data Buoy Center (UK), NCMR=National Center for Marine Research (Greece), KDI=Coastal Authorities (Denmark). H_s =significant wave height, H_m =maximum wave height, T_{02} =mean wave period, T_p =peak or dominant wave period, θ_w =mean wave direction. δt is the sampling rate in hours.

7.2 Observed wave statistics

| Station | min | mean | max | stdev |
|---------|-----|------|-----|-------|
| almag | 0.1 | 0.5 | 2.1 | 0.3 |
| oland | 0.1 | 0.7 | 2.8 | 0.4 |
| truba | 0.1 | 0.4 | 2.0 | 0.3 |
| 20044 | 0.2 | 0.7 | 2.3 | 0.4 |
| 24023 | 0.2 | 0.9 | 3.9 | 0.6 |
| 25077 | 0.2 | 0.9 | 3.8 | 0.6 |
| 25138 | 0.1 | 0.8 | 2.5 | 0.5 |
| 61001 | 0.2 | 0.9 | 3.0 | 0.6 |
| 61002 | 0.2 | 1.4 | 5.1 | 1.0 |
| 62001 | 0.8 | 2.5 | 6.7 | 1.0 |
| 62026 | 0.3 | 1.4 | 4.9 | 0.6 |
| 62029 | 1.0 | 3.1 | 7.9 | 1.3 |
| 62101 | 0.0 | 0.9 | 3.5 | 0.6 |
| 62103 | 0.3 | 1.3 | 4.6 | 0.7 |
| 62105 | 1.2 | 3.2 | 8.3 | 1.1 |
| 62106 | 1.1 | 2.9 | 8.9 | 1.0 |
| 62107 | 0.4 | 2.1 | 7.0 | 1.1 |
| 62108 | 1.1 | 3.2 | 8.8 | 1.3 |
| 62109 | 0.5 | 1.6 | 4.5 | 0.6 |
| 62117 | 0.4 | 1.4 | 4.0 | 0.6 |
| 62145 | 0.3 | 1.1 | 4.1 | 0.6 |
| 62163 | 1.0 | 2.7 | 7.5 | 1.2 |
| 62301 | 0.0 | 1.1 | 4.5 | 0.6 |
| 62303 | 0.4 | 1.8 | 5.7 | 1.0 |
| 62304 | 0.1 | 0.6 | 2.7 | 0.4 |
| 62305 | 0.1 | 0.9 | 5.4 | 0.7 |
| 62414 | 0.4 | 1.3 | 4.2 | 0.7 |
| 64045 | 1.1 | 2.9 | 9.2 | 1.1 |
| 64046 | 0.8 | 2.6 | 9.1 | 1.0 |
| athos | 0.1 | 0.5 | 1.5 | 0.3 |

Table 12. Observed wave height. The fraction of missing data is shown in Fig 4.

| Station | min | mean | max | stdev |
|---------|-----|------|-----|-------|
| almag | 2.5 | 3.6 | 5.2 | 0.5 |
| oland | 2.4 | 3.7 | 6.4 | 0.7 |
| truba | 2.1 | 3.2 | 5.3 | 0.5 |
| 20044 | 2.4 | 3.3 | 4.8 | 0.5 |
| 24023 | 2.3 | 3.9 | 6.9 | 0.8 |
| 25077 | 2.3 | 3.8 | 6.8 | 0.8 |
| 25138 | 2.1 | 3.4 | 5.3 | 0.7 |
| athos | 2.4 | 3.2 | 4.5 | 0.4 |

Table 13. Observed mean wave period

| Station | min | mean | max | stdev |
|---------|-----|------|------|-------|
| 20044 | 2.3 | 5.1 | 13.3 | 2.3 |
| 24023 | 1.9 | 7.2 | 16.7 | 3.0 |
| 25077 | 1.9 | 6.8 | 16.7 | 3.2 |
| 25138 | 1.9 | 5.1 | 16.7 | 2.5 |
| 61001 | 3.0 | 5.4 | 15.0 | 1.5 |
| 61002 | 3.0 | 5.4 | 13.0 | 1.5 |
| 62001 | 5.0 | 7.7 | 13.0 | 1.5 |
| 62026 | 3.0 | 5.1 | 8.0 | 0.8 |
| 62029 | 5.0 | 7.5 | 12.0 | 1.2 |
| 62101 | 3.0 | 4.2 | 7.0 | 0.9 |
| 62103 | 4.0 | 8.8 | 26.0 | 2.3 |
| 62105 | 2.0 | 7.5 | 13.0 | 1.2 |
| 62106 | 5.0 | 7.6 | 13.0 | 1.2 |
| 62107 | 5.0 | 8.7 | 17.0 | 1.5 |
| 62108 | 5.0 | 7.6 | 12.0 | 1.1 |
| 62109 | 4.0 | 5.2 | 8.0 | 0.8 |
| 62117 | 3.0 | 4.7 | 9.0 | 0.9 |
| 62145 | 3.0 | 4.2 | 7.0 | 0.8 |
| 62163 | 5.0 | 7.4 | 12.0 | 1.4 |
| 62301 | 3.0 | 4.7 | 10.0 | 1.2 |
| 62303 | 4.0 | 6.4 | 11.0 | 1.2 |
| 62304 | 2.0 | 6.2 | 8.0 | 0.7 |
| 62305 | 5.0 | 7.9 | 15.0 | 1.5 |
| 64045 | 3.0 | 7.4 | 12.0 | 1.1 |
| 64046 | 3.0 | 6.9 | 12.0 | 1.1 |
| athos | 2.0 | 4.0 | 6.2 | 1.0 |

Table 14. Observed dominant wave period

7.3 Wave height vs. wave period

The relation between significant wave height H_s and mean wave period T_{02} is shown on the diagrams, below, for those stations that record both quantities. At each station, there is a fair linear correlation between H_s and T_{02} , superimposed by a weak swell component.




Figure 12. Significant wave height vs. mean wave period

7.4 Wave forecast statistics

| Parameter Station | bias cm | % | rms cm | si | cc |
|----------------------|------------|-----|-----------|------|------|
| almag | 7 | 13 | 24 | 0.47 | 0.78 |
| oland | 7 | 10 | 26 | 0.37 | 0.86 |
| truba | 10 | 24 | 25 | 0.61 | 0.85 |
| 20044 | 6 | 8 | 27 | 0.38 | 0.80 |
| 24023 | 7 | 8 | 28 | 0.31 | 0.87 |
| 25077 | 10 | 12 | 29 | 0.32 | 0.89 |
| 25138 | -10 | -13 | 24 | 0.32 | 0.89 |
| 61001 | -16 | -18 | 38 | 0.44 | 0.81 |
| 61002 | -22 | -16 | 57 | 0.42 | 0.85 |
| 62001 | -37 | -15 | 55 | 0.22 | 0.92 |
| 62026 | -11 | -7 | 34 | 0.24 | 0.87 |
| 62029 | -17 | -5 | 52 | 0.17 | 0.93 |
| 62101 | 32 | 35 | 56 | 0.62 | 0.84 |
| 62103 | 52 | 40 | 72 | 0.56 | 0.87 |
| 62105 | -30 | -10 | 61 | 0.19 | 0.89 |
| 62106 | -21 | -7 | 54 | 0.19 | 0.89 |
| 62107 | 27 | 13 | 66 | 0.32 | 0.87 |
| 62108 | -33 | -10 | 65 | 0.20 | 0.90 |
| 62109 | -11 | -7 | 34 | 0.22 | 0.88 |
| 62117 | 17 | 12 | 42 | 0.30 | 0.84 |
| 62145 | 1 | 1 | 30 | 0.28 | 0.89 |
| 62163 | -16 | -6 | 44 | 0.16 | 0.94 |
| 62301 | -13 | -12 | 33 | 0.29 | 0.90 |
| 62303 | -10 | -6 | 45 | 0.25 | 0.91 |
| 62304 | 14 | 23 | 36 | 0.59 | 0.78 |
| 62305 | 21 | 23 | 43 | 0.47 | 0.88 |
| 62414 | -7 | -6 | 37 | 0.29 | 0.85 |
| 64045 | -14 | -5 | 54 | 0.19 | 0.90 |
| 64046 | -18 | -7 | 49 | 0.19 | 0.90 |
| athos | 4 | 7 | 28 | 0.54 | 0.53 |

Table 15. Predicted significant wave height

| Parameter Station | Obs m | peak | |
|----------------------|----------|------|-----|
| | | m | % |
| almag | 1.8 | -0.4 | -24 |
| oland | 2.3 | -0.4 | -15 |
| truba | 1.6 | 0.1 | 6 |
| 20044 | 1.5 | -0.1 | -5 |
| 24023 | 2.7 | 0.2 | 9 |
| 25077 | 2.8 | 0.0 | -1 |
| 25138 | 2.2 | -0.3 | -16 |
| 61001 | 2.5 | 0.0 | 0 |
| 61002 | 4.4 | -1.0 | -21 |
| 62001 | 5.9 | -1.0 | -17 |
| 62026 | 3.6 | -0.2 | -6 |
| 62029 | 7.6 | -0.9 | -11 |
| 62101 | 3.0 | 0.8 | 28 |
| 62103 | 3.9 | 1.0 | 24 |
| 62105 | 6.9 | -0.1 | -3 |
| 62106 | 6.4 | -0.3 | -5 |
| 62107 | 6.4 | -1.2 | -19 |
| 62108 | 7.8 | -1.0 | -12 |
| 62109 | 3.7 | 0.0 | 0 |
| 62117 | 1.7 | 0.1 | 6 |
| 62145 | 3.6 | -0.4 | -11 |
| 62163 | 6.7 | -0.6 | -9 |
| 62301 | 3.3 | 0.0 | -1 |
| 62303 | 5.2 | 0.1 | 1 |
| 62304 | 2.3 | -0.1 | -4 |
| 62305 | 3.9 | -0.1 | -1 |
| 62414 | 1.1 | 0.1 | 9 |
| 64045 | 6.4 | 0.2 | 0 |
| 64046 | 6.6 | -0.5 | -6 |
| athos | 1.2 | -0.2 | -16 |

Table 16. Average of top 5 wave events (peaks) and corresponding mean peak error (peak bias)

| Parameter Station | bias | | rms sec | si | cc |
|----------------------|------|----|------------|------|------|
| | sec | % | | | |
| almag | 0.4 | 11 | 0.6 | 0.17 | 0.57 |
| oland | 0.4 | 11 | 0.6 | 0.17 | 0.76 |
| truba | 0.6 | 19 | 0.9 | 0.27 | 0.44 |
| 20044 | 1.4 | 43 | 2.0 | 0.61 | 0.22 |
| 24023 | 1.9 | 49 | 2.3 | 0.58 | 0.60 |
| 25077 | 1.8 | 46 | 2.1 | 0.54 | 0.53 |
| 25138 | 1.7 | 51 | 2.2 | 0.67 | 0.25 |
| athos | 0.5 | 16 | 0.7 | 0.21 | 0.43 |

Table 17. Predicted mean wave period

| Parameter Station | bias sec | % | rms sec | si | cc |
|----------------------|-------------|-----|------------|------|------|
| 20044 | 0.3 | 7 | 2.1 | 0.41 | 0.57 |
| 24023 | 0.2 | 3 | 2.5 | 0.34 | 0.66 |
| 25077 | 0.4 | 5 | 2.6 | 0.39 | 0.64 |
| 25138 | 1.3 | 26 | 3.4 | 0.67 | 0.39 |
| 61001 | -0.2 | -5 | 1.9 | 0.35 | 0.27 |
| 61002 | -0.1 | -1 | 1.4 | 0.27 | 0.54 |
| 62001 | 2.9 | 37 | 3.4 | 0.44 | 0.63 |
| 62026 | 1.3 | 26 | 2.3 | 0.46 | 0.39 |
| 62029 | 3.0 | 40 | 3.4 | 0.45 | 0.67 |
| 62101 | 3.4 | 81 | 4.3 | 1.04 | 0.42 |
| 62103 | 0.6 | 7 | 2.5 | 0.29 | 0.44 |
| 62105 | 2.9 | 38 | 3.2 | 0.43 | 0.67 |
| 62106 | 2.8 | 37 | 3.2 | 0.43 | 0.68 |
| 62107 | 1.4 | 16 | 2.1 | 0.24 | 0.71 |
| 62108 | 2.5 | 33 | 2.9 | 0.38 | 0.64 |
| 62109 | 1.5 | 28 | 2.4 | 0.46 | 0.24 |
| 62117 | 3.0 | 63 | 3.8 | 0.81 | 0.17 |
| 62145 | 1.4 | 33 | 1.8 | 0.43 | 0.60 |
| 62163 | 3.1 | 42 | 3.4 | 0.46 | 0.73 |
| 62301 | 3.0 | 63 | 4.0 | 0.84 | 0.51 |
| 62303 | 2.5 | 39 | 3.2 | 0.50 | 0.61 |
| 62304 | -0.9 | -15 | 1.8 | 0.30 | 0.39 |
| 62305 | -1.6 | -20 | 2.9 | 0.36 | 0.24 |
| 64045 | 2.8 | 38 | 3.3 | 0.44 | 0.67 |
| 64046 | 2.9 | 42 | 3.4 | 0.50 | 0.59 |
| athos | 0.1 | 1 | 0.9 | 0.24 | 0.37 |

Table 18. Predicted dominant wave period

| Parameter Station | bias deg | std deg | cc |
|----------------------|-------------|------------|------|
| 20044 | 14 | 58 | 0.84 |
| 24023 | -1 | 41 | 0.80 |
| 25077 | 1 | 38 | 0.87 |
| 25138 | 6 | 49 | 0.85 |
| athos | 14 | 49 | 0.89 |

Table 19. Predicted mean wave direction

7.5 Significant wave height station plots

The following pages show significant wave height error statistics for each station.




Figure 13. Significant wave height: Almagrundet




Figure 14. Significant wave height: Øland




Figure 15. Significant wave height: Trubaduren




Figure 16. Significant wave height: 20044




Figure 17. Significant wave height: 24023. The zig-zag curves are caused by irregular 3h sampling.




Figure 18. Significant wave height: 25077. The zig-zag curves are caused by irregular 3h sampling.




Figure 19. Significant wave height: 25138




Figure 20. Significant wave height: 61001




Figure 21. Significant wave height: 61002




Figure 22. Significant wave height: 62001




Figure 23. Significant wave height: 62026




Figure 24. Significant wave height: 62029




Figure 25. Significant wave height: 62101. The sampling accuracy is 0.5m




Figure 26. Significant wave height: 62103




Figure 27. Significant wave height: 62105. A single event (June 17th) is grossly overpredicted on short range due to error in the wind forecast.




Figure 28. Significant wave height: 62106. A single event (June 17th) is grossly overpredicted on short range due to error in the wind forecast.




Figure 29. Significant wave height: 62107




Figure 30. Significant wave height: 62108




Figure 31. Significant wave height: 62109




Figure 32. Significant wave height: 62117. The zig-zag curves are caused by irregular 3h sampling.




Figure 33. Significant wave height: 62145




Figure 34. Significant wave height: 62163




Figure 35. Significant wave height: 62301. The sampling accuracy is 0.5m




Figure 36. Significant wave height: 62303




Figure 37. Significant wave height: 62304




Figure 38. Significant wave height: 62305




Figure 39. Significant wave height: 62414. The zig-zag curves are caused by irregular 3h sampling.




Figure 40. Significant wave height: 64045. A single event (June 17th) is grossly overpredicted on short range due to error in the wind forecast.




Figure 41. Significant wave height: 64046




Figure 42. Significant wave height: Athos

7.6 Mean wave period station plots

The following pages show mean wave period error statistics for each station. Only stations with reasonable statistics are included.




Figure 43. Mean wave period: Almagrundet




Figure 44. Mean wave period: Øland




Figure 45. Mean wave period: Trubaduren




Figure 46. Mean wave period: Athos

7.7 Dominant wave period station plots

The following pages show dominant wave period error statistics for each station. Only stations with reasonable statistics are included.




Figure 47. Dominant wave period: 61001




Figure 48. Dominant wave period: 61002




Figure 49. Dominant wave period: Athos

References

- [1] G. Komen et al. *Dynamics and Modelling of Ocean Waves*. Cambridge University Press, 1994.
- [2] The SWAMP group. *Ocean Wave Modelling*. Plenum Press, New York, 1985.
- [3] H. Günther, S. Hasselmann, and P. Janssen. Wamodel cycle4 (revised version). Technical Report 4, Deutches Klimarechnenzentrum, 1992.
- [4] J. W. Nielsen. Verification of DMI wave forecasts: 1st quarter of 2002. Technical Report 02-20, Danish Meteorologiacl Institute, 2002.
- [5] J. W. Nielsen, J. B. Jørgensen, and J. She. Verification of wave forecasts: DMI-WAM nov-dec 2001. Technical Report 02-18, Danish Meteorologiacl Institute, 2002.
- [6] J. She. HIRLAM-WAM quality assessment on winds and waves in the North Sea. Scientific Report 00-27, Danish Meteorological Institute, 2000.
- [7] J. She and J.W. Nielsen. Operational wave forecasts in Baltic and North Sea. Scientific Report 99-07, Danish Meteorological Institute, 1999.

List of Tables

| | | |
|----|---|----|
| 1 | Key numbers | 3 |
| 2 | DMI-WAM operational set-up | 5 |
| 3 | DMI-WAM wave parameters | 6 |
| 4 | Wave data providers. | 6 |
| 5 | Wave stations in each domain | 7 |
| 6 | Significant wave height results | 10 |
| 7 | Extreme wave height results | 13 |
| 8 | Mean wave period results | 14 |
| 9 | Dominant wave period results | 15 |
| 10 | Mean wave direction results | 16 |
| 11 | Wave stations and locations | 18 |
| 12 | Observed wave height | 19 |
| 13 | Observed mean wave period | 20 |
| 14 | Observed dominant wave period | 20 |
| 15 | Predicted significant wave height | 22 |
| 16 | Predicted peak waves and errors | 23 |
| 17 | Predicted mean wave period | 23 |
| 18 | Predicted dominant wave period | 24 |
| 19 | Predicted mean wave direction | 24 |

List of Figures

| | | |
|----|---|----|
| 1 | DMI wave models | 4 |
| 2 | DMI Hirlam | 5 |
| 3 | Wave recorders | 7 |
| 4 | Missing data | 8 |
| 5 | Significant wave height: short range (1-6h) scatter diagram | 10 |
| 6 | Significant wave height: scatter index | 11 |
| 7 | Significant wave height | 12 |
| 8 | Extreme wave height | 13 |
| 9 | Mean wave period | 14 |
| 10 | Dominant wave period | 15 |
| 11 | Mean wave direction | 16 |
| 12 | Significant wave height vs. mean wave period | 21 |
| 13 | Significant wave height: Almagrundet | 26 |
| 14 | Significant wave height: Øland | 27 |
| 15 | Significant wave height: Trubaduren | 28 |
| 16 | Significant wave height: 20044 | 29 |
| 17 | Significant wave height: 24023 | 30 |
| 18 | Significant wave height: 25077 | 31 |
| 19 | Significant wave height: 25138 | 32 |
| 20 | Significant wave height: 61001 | 33 |
| 21 | Significant wave height: 61002 | 34 |
| 22 | Significant wave height: 62001 | 35 |
| 23 | Significant wave height: 62026 | 36 |
| 24 | Significant wave height: 62029 | 37 |
| 25 | Significant wave height: 62101 | 38 |

| | | |
|----|--|----|
| 26 | Significant wave height: 62103 | 39 |
| 27 | Significant wave height: 62105 | 40 |
| 28 | Significant wave height: 62106 | 41 |
| 29 | Significant wave height: 62107 | 42 |
| 30 | Significant wave height: 62108 | 43 |
| 31 | Significant wave height: 62109 | 44 |
| 32 | Significant wave height: 62117 | 45 |
| 33 | Significant wave height: 62145 | 46 |
| 34 | Significant wave height: 62163 | 47 |
| 35 | Significant wave height: 62301 | 48 |
| 36 | Significant wave height: 62303 | 49 |
| 37 | Significant wave height: 62304 | 50 |
| 38 | Significant wave height: 62305 | 51 |
| 39 | Significant wave height: 62414 | 52 |
| 40 | Significant wave height: 64045 | 53 |
| 41 | Significant wave height: 64046 | 54 |
| 42 | Significant wave height: Athos | 55 |
| 43 | Mean wave period: Almagrundet | 57 |
| 44 | Mean wave period: Øland | 58 |
| 45 | Mean wave period: Trubaduren | 59 |
| 46 | Mean wave period: Athos | 60 |
| 47 | Dominant wave period: 61001 | 62 |
| 48 | Dominant wave period: 61002 | 63 |
| 49 | Dominant wave period: Athos | 64 |